

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

CLAIMS

1. A method of producing a hybrid DNA molecule having a sense strand and an anti-sense strand and in which, reading in the 5' to 3' direction, the sense strand has the sequences x_1, x_2, \dots, x_n , where n is greater than or equal to 3, the method comprising the steps of

- (1) providing in a single reaction mixture
 - (a) the sequences x_1, x_2, \dots, x_n and their complementary sequences x_1', x_2', \dots, x_n' , to be assembled into the hybrid molecule,
 - (b) for each pair of complementary sequences defined in (a) a respective pair of PCR primers each having a priming sequence and which are such that the primers hybridising to the 3' ends of any two sequences (x_i, x_{i+1}'), where i is 1 to $(n-1)$, have specifically complementary linker sequences
- (2) effecting a first stage PCR reaction in which those primers provided with linker sequences are present in limiting concentrations, and
- (3) effecting a second stage PCR reaction using a single pair of primers one of which provides the 5'-end of the sense strand and other of which provides the 3'-end of the anti-sense strand of the required hybrid molecule

whereby said hybrid molecule is generated.

2. A method as claimed in claim 1 wherein the polymerising enzyme adds a 3' adenosine overhang to an extended strand and those primers incorporating linker

WO 99/64624

PCT/GB99/01691

29

sequences have their priming sequences connected to their respective linker sequences via an adenine residue.

3. A method as claimed in claim 2 wherein the polymerising enzyme is *Taq*.
4. A method as claimed in any one of claims 1 to 3 wherein the annealing temperature (T_m) of the linker sequences is greater than that of the priming sequences to the x and x' sequences.
5. A method as claimed in claim 4 wherein the annealing temperature of the linker sequences is 2 to 5°C greater than that of the priming sequences to the x and x' sequences.
6. A method as claimed in any one of claims 1 to 5 wherein the linker sequences do not have intrinsic secondary structure.
7. A method as claimed in any one of claims 1 to 6 wherein between the first and second stage PCR reactions the reaction mixture is frozen to deactivate residual PCR activity.
8. A method as claimed in any one of claims 1 to 6 wherein between the first and second stage PCR reactions the reaction mixture is treated with an exonuclease I to digest single stranded molecules.
9. A method as claimed in any one of claims 1 to 8 wherein each of the first and second stage PCR reactions utilise a thermally activated polymerase.
10. A method of mutation analysis wherein the analysis is effected on a DNA hybrid molecule produced in accordance with the method of any one of claims 1 to 9.

WO 99/64524

PCT/GB99/01691

30

11. A set of primers incorporating the following sequences.

5'tcatattgcccgtgcattgcc-a-3'

5'ggcaatgcagcggctaataatga-a-3'

5'agccatgacccaactcctgt-a-3'

5'acaggagtttggtagtggt-a-3'

5'tgtctcactgaaccgcctacct-a-3'

5'aggtaggcagggttcagttagaca-a-3'

5'cctcat taccggctgtcagactg-a-3'

5'cagctgacagccggtaatgagg-a-3'

12. A method of producing a hybrid DNA molecule having a sense strand and an anti-sense strand and in which, reading in the 5' to 3' direction, the sense strand has the sequences x_1, x_2, \dots, x_n , where n is greater than or equal to 3, the method comprising the steps of

- (a) providing in a single reaction mixture
 - (a) the sequences x_1, x_2, \dots, x_n and their complementary sequences x'_1, x'_2, \dots, x'_n , to be assembled into the hybrid molecule,
 - (b) for each pair of complementary sequences defined in (a) a respective pair of PCR primers each having a priming sequence and which are such that the primers for the 3' ends of any two sequences $(x_i, x'_{(i+1)})$, where i is 1 to $(n-1)$, have specifically complementary linker sequences connected to their respective priming sequences via an adenine residue, and

WO 99/64624

PCT/GB99/01691

31

- (2) effecting a PCR reaction using a polymerase which adds a 3' adenine overhang to the end of an extended strand